

I. Healthcare Personnel (HCP) Vaccination Requirements:

| Vaccine | Requirements |
|--------------------------------|---|
| Hepatitis B | Give 3-dose series (dose #1 now, #2 in 1 month, #3 approximately 5 months after #2). Give IM. Obtain anti-HBs serologic testing 1-2 months after dose #3. |
| Influenza | Give 1 dose of influenza vaccine annually. Give inactivated injectable influenza vaccine intramuscularly or live attenuated influenza vaccine (LAIV) intranasally. By November 30 th . If HCP does not get vaccine they must wear a N95 mask for the duration of their shift. |
| MMR | For healthcare personnel (HCP) born in 1957 or later without serologic evidence of immunity or prior vaccination, give 2 doses of MMR, 4 weeks apart. It is recommended that HCP born prior to 1957 who do not have laboratory evidence of disease or immunity to measles, mumps and/or rubella, receive 2 doses of MMR vaccine. Give SC. |
| Varicella (chickenpox) | For HCP who have no serologic proof of immunity, prior vaccination, or history of varicella disease, give 2 doses of varicella vaccine, 4 weeks apart. Give SC. |
| Tetanus, diphtheria, pertussis | A tetanus immunization without the diphtheria/pertussis components is not acceptable. Give a one-time dose of Tdap as soon as feasible to all HCP who have not received Tdap previously. Give Tdap boosters every 10 years thereafter. Give IM. |
| Meningococcal | Give 1 dose to microbiologists who are routinely exposed to isolates of <i>N. meningitidis</i> . Give IM or SC. |